

SUBSTITUTE SHEET

RECEPTOR BASED ANTAGONISTS AND METHODS OF MAKING AND USING

This application claims priority of U.S. S.N. 09/313,942 filed May 19, 1999, which claims priority of Provisional application No. 60/101,858 filed September 25, 1998.

5 Throughout this application various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application.

BACKGROUND OF THE INVENTION

10 Although discovered for varying biological activities, ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), oncostatin M (OSM) and interleukin-6 (IL-6) comprise a defined family of cytokines (referred to herein as the "CNTF family" of cytokines). These cytokines are grouped together
15 because of their distant structural similarities [Bazan, J. Neuron 7: 197-208 (1991); Rose and Bruce, Proc. Natl. Acad. Sci. USA 88: 8641-8645 (1991)], and, perhaps more importantly, because they share " β " signal-transducing receptor components [Baumann, et al., J. Biol. Chem. 265:19853-19862 (1993); Davis, et al., Science 260: 1805-1808 (1993); Gearing et al., Science 255:1434-1437 (1992); Ip
20 et al., Cell 69: 1121-1132 (1992); Stahl, et al., J. Biol. Chem. 268: 7628-7631 (1993); Stahl and Yancopoulos, Cell 74: 587-590 (1993)]. Receptor activation by this family of cytokines results from either homo- or hetero-dimerization of these β components [Davis, et al. Science 260: 1805-1808 (1993), Murakami, et al., Science 260: 1808-1810 (1993); Stahl and Yancopoulos, Cell 74: 587-590 (1993)].
25 IL-6 receptor activation requires homodimerization of gp130 [Murakami, et al. Science 260: 1808-1810 (1993), Hibi, et al., Cell 63: 1149-1157 (1990)], a protein initially identified as the IL-6 signal transducer [Hibi, et al., Cell 63: 1149-1157